INTERNATIONAL JOURNAL OF PHARMACEUTICS

VOL. 98 NO. 1–3 CONTENTS 31 AUGUST 1993

Cited in: Current Contents (Life Sciences; Clinical Practice); International Pharmaceutical Abstracts; Biological Abstracts; Chemical Abstracts; Excerpta Medica EMBASE; PASCAL M; Science Citation Index; Current Awareness in Biological Sciences

Announcement from the Publisher v

Research Papers

QSARs of some novel antibacterial benzimidazoles, benzoxazoles, and oxazolopyridines against an enteric gram-negative rod; *K. pneumoniae*
I. Yalçin and E. Şener (Turkey) 1

Lymph node targeting and pharmacokinetics of [^{3}H]methotrexate-encapsulated neutral large unilamellar vesicles and immunoliposomes
C.-K. Kim, Y.J. Choi, S.-J. Lim, M.G. Lee, S.H. Lee and S.J. Hwang (South Korea) 9

Lipid vesicles loaded with thymopentin: characterization and in vitro activity on tumoral cells
A.M. Panico, R. Pignatello, S. Mazzone, V. Cardile, M. Bindoni, F. Cordopatri, A. Villari, N. Micali and G. Mazzone (Italy) 19

Dissolution kinetics of piroxicam in solid dispersions with polyethylene glycol 4000
M. Fernández, M.V. Margarit, I.C. Rodríguez and A. Cerezo (Spain) 29

In vitro percutaneous absorption of naproxen from gels using a double-layer artificial membrane
M.D. Contreras Claramonte, A. Parera Vialard and F.G. Vilchez (Spain) 37

Ultrasonic nebulisation of pentamidine isethionate
K.M.G. Taylor and C. Hoare (UK) 45

Theophylline inhibits the elimination of flumazenil in rabbits
T.A. Najjar, M.I. Al-Hassan and R.M.A. Khan (Saudi Arabia) 51

Probing the mechanisms of swelling of hydroxypropylmethylcellulose matrices
E. Papadimitriou, G. Buckton and M. Efentakis (UK) 57

Radiolabelling of polymer microspheres for scintigraphic investigations by neutron activation. 2. Effects of irradiation on the properties of Eudragit RS-sulphasalazine microspheres
P.J. Watts, C.G. Wilson, M.C. Davies and C.D. Melia (UK) 63

Radiolabelling of polymer microspheres for scintigraphic investigations by neutron activation. 3. Changes in the physical properties of Eudragit RS-sulphapyridine microspheres from incorporating samarium oxide
P.J. Watts, C.G. Wilson, M.C. Davies and C.D. Melia (UK) 75

pH-dependent transepithelial transport of cephalexin in rabbit intestinal mucosa
I.J. Hidalgo, F.M. Ryan, G.J. Marks and P.L. Smith (USA) 83

In vitro percutaneous absorption evaluation of phenobarbital through hairless mouse, adult and premature human skin
F.P. Bonina, L. Montenegro, G. Micali, D.P. West, P. Palicharla and R.L. Koch (Italy, USA) 93

Solubility and in vitro percutaneous absorption of tetracaine from solvents of propylene glycol and saline
K.J. Miller II, Y.K. Rao, S.R. Goodwin, G.B. Westermann-Clark and D.O. Shah (USA) 101

Percutaneous absorption of terodiline and the membrane-controlled transdermal therapeutic system
T. Ogiso, M. Iwaki, T. Tanino, M. Yatomi and C. Tsujimoto (Japan) 113

Conjugate of mitomycin C with *N*-succinyl-chitosan: In vitro drug release properties, toxicity and antitumor activity
Y. Song, H. Onishi and T. Nagai (Japan) 121

Penetration of spin-labeled retinoic acid from liposomal preparations into the skin of SKH1 hairless mice. Measurement by EPR tomography
C. Michel, N. Groth, T. Herrling, P. Rudolph, J. Fuchs, J. Kreuter and H.-J. Freisleben (Germany) 131

Improved stability of the human epidermal membrane during successive permeability experiments
K.D. Peck, A.H. Ghanem, W.I. Higuchi and V. Srinivasan (USA) 141

Stability of antibody-bearing liposomes containing dideoxyinosine triphosphate
G.V. Betageri and L.S. Burrell (USA) 149

Influence of manufacturing parameters on the size characteristics and the release profiles of nifedipine from poly(DL-lactide-co-glycolide) microspheres
P. Sansdrap and A.J. Moës (Belgium) 157

(Contents continued on p. 250)

0378-5173(19930831)98:1/3;1-6

(Contents continued from back cover)

Studies on the gastrointestinal absorption of phosphocitrate, a powerful controller of hydroxyapatite formation
C.M. Cooper and J.D. Sallis (Australia) 165
Azone® and the formation of reversed mono- and bicontinuous lipid-water phases
J. Engblom and S. Engström (Sweden) 173
Use of chitosan and chitosan malate as an excipient in wet granulation of three water soluble drugs
I. Henriksen, Ø. Skaugrud and J. Karlsen (Norway) 181
The mechanism of phenytoin crystal growth
G.L. Zipp and N. Rodríguez-Hornedo (USA) 189
Fluorescence analysis with diphenylhexatriene and its ionic derivatives of the fluidity of liposomes constituted from stratum corneum lipids: Contribution of each lipid component and effects of long-chain unsaturated fatty acids
S. Kitagawa, M. Sawada and H. Hirata (Japan) 203
Evaluation of reaction order. Software in pharmaceutics: III
S.A. Sande and J. Karlsen (Norway) 209
Preparation, characterization and pharmaceutical application of linear dextrins: IV. Drug release from capsules and tablets containing amylodextrin
G.H.P. Te Wierik, A.C. Eissens and C.F. Lerk (The Netherlands) 219
The effect of cyclodextrins on the solubility and stability of medroxyprogesterone acetate and megestrol acetate in aqueous solution
T. Loftsson, J. Baldvinsdóttir and A.M. Sigurdardóttir (Iceland) 225

Notes
Effects of tetracycline on the 2 h urinary and salivary excretion of metronidazole
K.E. Ibrahim, E.I. A/Karim, B.M. Ahmed and I.M. Nour (Sudan) 231
Polycarbophil as a controlled release matrix
E.A. Hosny (Saudi Arabia) 235
Increase in temperature enhances solubility of drugs in aqueous solutions of hydroxypropylcyclodextrins
T. Hoshino, K. Uekama and J. Pitha (USA, Japan) 239

Author Index 243

Noticeboard 247